



AMENDMENTS TO THE CLAIMS


1. (Currently Amended) A photothermographic material comprising, on one side of a support, a photosensitive silver halide, a non-photosensitive silver salt of an organic acid, a reducing agent for silver ions and a binder, which is characterized by containing at least one or more e-polyphenol compounds, phenol compound as the reducing agent and one or more compounds wherein all of said one or more compounds satisfy
at least one compound having the following requirements A and B-A: the compound has a hydrogen bond formation rate constant K_f that is 20-4000, B: the compound and which is represented by the following formula (II), (III), (IV) or (V) or the compound has a phosphoryl group (III) or (IV):




—(H)—



(III)



(IV)



—(V)—

wherein:


in the formula (II), R^{21} and R^{22} independently represent an alkyl group, and R^{23} represents an alkyl group, an aryl group or a heterocyclic group, and two or more of R^{21} , R^{22} and R^{23} may be taken together to form a ring;

in the formula (III), R^{31} and R^{32} independently represent an alkyl group, an aryl group, an aryl group or a heterocyclic group, and R^{31} and R^{32} may be taken together to form a ring;

and in the formula (IV), R^{41} and R^{42} independently represent an alkyl group, an aryl group or a heterocyclic group, R^{43} represents an alkyl group, an aryl group, a heterocyclic group or $\text{N-}(\text{R}^{44})(\text{R}^{45})$ where R^{44} and R^{45} independently represent an alkyl group, an aryl group or a heterocyclic group, and two or more of R^{41} , R^{42} , R^{43} , R^{44} and R^{45} may be taken together to form a ring, ring.

and in the formula (V), R^{51} , R^{52} , R^{53} , R^{54} and R^{55} independently represent a hydrogen atom or a substituent and two or more of R^{51} , R^{52} , R^{53} , R^{54} and R^{55} may be taken together to form a ring.

2. (Currently Amended) The photothermographic material according to claim 1, wherein the phenol compound is at least one of the o-polyphenol compounds is compound represented by the following formula I



(I)

wherein R^2 , R^4 , R^5 , and R^7 are hydrogen atoms, R^1 and R^8 represent an alkyl group and R^3 and R^6 represent an alkyl group, and L represents a group $-CHR^9-$ where R^9 represents a hydrogen atom, a methyl group, an ethyl group, an isopropyl group, an n-propyl group, a heptyl group, a 1-ethylpentyl group, and an undecyl group.

3. (Original) The photothermographic material according to claim 1 or 2, wherein the hydrogen bond formation rate constant Kf is 70 to 4000.

4. (Currently Amended) The photothermographic material according to claim 1 or 2, wherein the hydrogen bound bond formation rate constant Kf is 100-4000.

5. (Currently Amended) The photothermographic material according to claim 1 or 2, wherein the hydrogen bound bond formation rate constant Kf is 250-2000.

6. (Cancelled)

7. (Original) The photothermographic material according to claim 1 or 2, wherein the compound of the requirement B is represented by the formula (III).

8. (Original) The photothermographic material according to claim 1 or 2, wherein the compound of the requirement B is represented by the formula (IV).

9. (Cancelled)

10. (Currently Amended) The photothermographic material according to claim 1 or 2, wherein the amount of the ~~o-polyphenol phenol~~ compound is 0.01-40 g/m².

11. (Currently Amended) The photothermographic material according to claim 1 or 2, wherein the amount of ~~all of~~ said at least one ~~or more~~ compound ~~satisfy the requirement A and B~~ is 0.01-40g/m².